


INDOL-6 SULFONAMIDE DERIVATIVES, THEIR PREPARATION AND THEIR USE 5-HT-6 ~~ALS~~
MODULATORS

AS

- 5 The present invention relates to new sulfonamide derivatives, of general formula (Ia, Ib, Ic),



(Ia, Ib, Ic),

- 10 optionally in form of one of their stereoisomers, preferably enantiomers or diastereomers, a racemate, or in form of a mixture of at least two of their stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or their salts, preferably the corresponding, physiologically acceptable salts, or corresponding solvates; to the processes for their preparation, to their
15 application in medicaments in human and/or veterinary therapeutics, and to pharmaceutical compositions containing them.

- The new compounds of the present invention can be used in the pharmaceutical industry as intermediates and for the manufacture of
20 medicaments.

- The superfamily of serotonin receptors (5-HT) comprises 7 classes (5-HT₁-5-HT₇), which cover 14 human subclasses [D. Hoyer, et al., *Neuropharmacology*, 1997, 36, 419]. The 5-HT₆ receptor has been the last
25 serotonin receptor identified by molecular cloning in rats [F.J. Monsma, et al., *Mol. Pharmacol.*, 1993, 43, 320; M. Ruat, et al., *Biochem. Biophys. Res. Commun.*, 1993, 193, 268] as well as in humans [R. Kohen, et al., *J.*

Neurochem., 1996, 66, 47]. The compounds with an affinity for the 5-HT₆ receptor are useful in treating different disorders of the Central Nervous System and of the Gastrointestinal system, as well as the irritable bowel syndrome. The compounds with an affinity for the 5-HT₆ receptor are useful for treating anxiety, depression and cognitive memory disorders [M. Yoshioka, et al., *Ann. NY Acad. Sci.*, 1998, 861, 244; A. Bourson, et al., *Br. J. Pharmacol.*, 1998, 125, 1562; D.C. Rogers, et al., *Br. J. Pharmacol. Suppl.*, 1999, 127, 22P; A. Bourson, et al., *J. Pharmacol. Exp. Ther.*, 1995, 274, 173; A.J. Sleight, et al., *Behav. Brain Res.*, 1996, 73, 245; T.A. Branchek, et al., *Annu. Rev. Pharmacol. Toxicol.*, 2000, 40, 319; C. Routledge, et al., *Br. J. Pharmacol.*, 2000, 130, 1606]. It has been shown that the typical and atypical antipsychotics for treating schizophrenia have a high affinity for the 5-HT₆ receptors [B.L. Roth, et al., *J. Pharmacol. Exp. Ther.*, 1994, 268, 1403; C.E. Glatt, et al., *Mol. Med.*, 1995, 1, 398; F.J. Mosma, et al., *Mol. Pharmacol.*, 1993, 43, 320; T. Shinkai, et al., *Am. J. Med. Genet.*, 1999, 88, 120]. The compounds with an affinity for the 5-HT₆ receptor are useful for treating infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) [W.D. Hirst, et al., *Br. J. Pharmacol.*, 2000, 130, 1597; C. Gérard, et al., *Brain Research*, 1997, 746, 207; M.R. Pranzatelli, *Drugs of Today*, 1997, 33, 379].

Patent application WO 01/32646 discloses sulfonamides derived from bicycles, whereby each of the rings is 6-membered, aromatic or heteroaromatic rings with 5-HT₆ receptor antagonist activity.

Patent application EP 0 733 628 discloses sulfonamides derived from indole with 5-HT_{1F} receptor antagonist activity, useful for the treatment of migraines.

Furthermore, it has been shown that the 5-HT₆ receptor plays a role in the ingestion of food [Neuropharmacology, 41, 2001, 210-219].

Eating disorders, particularly obesity, are a serious and increasingly frequent threat for the health of humans from all ages, since these diseases increase the risk of developing other serious and even mortal diseases, preferably diabetes and coronary artery diseases.


Therefore, an object of the present invention was to provide new compounds, particularly suitable as active substances in medicaments, preferably in medicaments for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder), and other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in animals, preferably in mammals, more preferably in humans.

It has been found that the indol-6-yl sulfonamide compounds of general formulas (Ia, Ib, Ic) described below show an affinity for the 5-HT₆ receptor. These compounds are therefore suitable for the manufacture of a medicament for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis

-4-

or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) and other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in animals, preferably in mammals, more preferably in humans.

- 5 Thus, one aspect of the present invention are compounds of general formula (Ia),



(Ia)

wherein

- 10 R¹ represents a -NR⁸R⁹ radical or a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing cycloaliphatic radical, which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system,
- 15

- R², R³, R⁴, R⁵ and R⁷, identical or different, each represent hydrogen, halogen, nitro, alkoxy, cyano, a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical, or an optionally at least mono-substituted phenyl or optionally at least mono-substituted heteroaryl radical,
- 20

R⁶ represents hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical,

R^8 and R^9 , identical or different, each represent hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical,

5 with the proviso that R^8 and R^9 are not hydrogen at the same time, and if one of them, R^8 or R^9 , is a saturated or unsaturated, linear or branched, optionally at least mono-substituted C_1 - C_4 aliphatic radical, the other one is a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical with at least five carbon atoms,

10

or

R^8 and R^9 , together with the bridging nitrogen atom, form a saturated or unsaturated, optionally at least mono-substituted heterocyclic ring, which may
15 contain at least one further heteroatom as a ring member and/or which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system,

20 A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, which may be bonded via an optionally at least mono-substituted alkylene, alkenylene or alkynylene group and/or which may contain at least one heteroatom as a ring member in one or more of its rings,

25 and


n is 0, 1, 2, 3 or 4;

optionally in form of one of its stereoisomers, preferably enantiomers or
30 diastereomers, a racemate or in form of a mixture of at least two of its stereoisomers, preferably enantiomers and/or diastereomers, in any mixing ratio, or a salt thereof, preferably a corresponding, physiologically acceptable

-6-

salt thereof, or a corresponding solvate thereof.

Another aspect of the present invention are compounds of general formula (Ib)



(Ib)

5

wherein

R^1 is a $-NR^8R^9$ radical,

10 R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen, halogen, nitro, alkoxy, cyano, a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical, or an optionally at least mono-substituted phenyl or optionally at least mono-substituted heteroaryl radical,

15 R^6 represents hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical,

20 R^8 and R^9 , identical or different, each represent hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted C_1 - C_4 aliphatic radical,


A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, which may be bonded via an optionally at least mono-substituted alkylene, alkenylene or alkynylene group and/or which may contain
25 at least one heteroatom as a ring member in one or more of its rings,

n is 0, 1, 2, 3 or 4;

-7-

optionally in form of one of its stereoisomers, preferably enantiomers or diastereomers, a racemate or in form of a mixture of at least two of its stereoisomers, preferably enantiomers and/or diastereomers, in any mixing
 5 ratio, or a salt thereof, preferably a corresponding, physiologically acceptable salt thereof, or a corresponding solvate thereof.

Yet, another aspect of the present invention are compounds of general formula (Ic),



10

(Ic),

wherein

15 R^1 represents a $-NR^8R^9$ radical or a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing cycloaliphatic radical, which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one
 20 heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system,

R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen, halogen, nitro, alkoxy, cyano, a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical, or an optionally at least mono-
 25 substituted phenyl or optionally at least mono-substituted heteroaryl radical,

-8-

R^6 represents hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical,

5 R^8 and R^9 , identical or different, each represent hydrogen or a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical,

or

10 R^8 and R^9 , together with the bridging nitrogen atom, form a saturated or unsaturated, optionally at least mono-substituted heterocyclic ring, which may contain at least one further heteroatom as a ring member and/or which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing
15 mono- or bicyclic cycloaliphatic ring system,

A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, which may be bonded via an optionally at least mono-substituted alkylene, alkenylene or alkynylene group and/or which may contain
20 at least one heteroatom as a ring member in one or more of its rings,

and

n is 0, 1, 2, 3 or 4;
25

optionally in form of one of its stereoisomers, preferably enantiomers or diastereomers, a racemate or in form of a mixture of at least two of its stereoisomers, preferably enantiomers and/or diastereomers, in any mixing ratio, or a salt thereof, preferably a corresponding, physiologically acceptable
30 salt thereof, or a corresponding solvate thereof.

If one or more of the moieties R^2 to R^9 represents a saturated or unsaturated aliphatic radical, that is, an alkyl, alkenyl or alkynyl radical, which is substituted by one or more substituents, each one of these substituents may preferably be chosen, unless otherwise defined, from the group consisting of hydroxy, fluorine, chlorine, bromine and trifluoromethyl.

If R^1 is a saturated or unsaturated, optionally at least one heteroatom as a ring member containing cycloaliphatic radical, which is substituted by one or more substituents and/or is condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system, each of these substituents may, unless otherwise defined, preferably be chosen from the group consisting of hydroxy, fluorine, chlorine, bromine, linear or branched C_1 - C_6 alkyl, linear or branched C_1 - C_6 alkoxy, linear or branched C_1 - C_6 perfluoroalkyl, linear or branched C_1 - C_6 perfluoroalkoxy and benzyl, more preferably chosen from the group consisting of linear or branched C_1 - C_6 alkyl and benzyl.

The heteroatoms of said cycloaliphatic radical and/or of the mono- or bicyclic cycloaliphatic ring may, independently from one another, be chosen preferably from the group consisting of nitrogen, sulphur and oxygen, more preferably nitrogen is chosen as a heteroatom.

Said cycloaliphatic radical may contain 0, 1, 2 or 3 heteroatoms chosen from the above mentioned group, preferably it contains 0, 1 or 2 heteroatoms chosen from the above mentioned group.

If R^8 and R^9 together with the bridging nitrogen atom form a saturated or unsaturated, optionally at least mono-substituted heterocyclic ring, which may contain at least one further heteroatom as a ring member and/or which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally one heteroatom as a ring member containing mono- or

bicyclic ring system, each one of these substituents may, unless otherwise defined, preferably be chosen from the group consisting of hydroxy, fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ perfluoroalkyl, linear or branched C₁-C₆ perfluoroalkoxy and benzyl, more preferably from the group consisting of linear or branched C₁-C₆ alkyl and benzyl.

If the heterocyclic ring contains one or more additional heteroatoms, and/or if one or both mono- or bicyclic rings contain one or more additional heteroatoms, these heteroatoms may, independently from one another, preferably be chosen from the group consisting of nitrogen, sulphur and oxygen, more preferably nitrogen is chosen as a heteroatom.

Said heterocyclic ring may contain 0, 1, 2 or 3 additional heteroatoms chosen from the above mentioned group, preferably it contains 0 or 1 heteroatoms chosen from the above mentioned group.

If A is an optionally at least one heteroatom as a ring member containing mono- or polycyclic aromatic ring system, which may be bonded via an optionally at least mono-substituted alkylene, alkenylene or alkynylene group and which may be substituted by one or more substituents, each of these substituents may preferably be chosen, unless otherwise defined, from the group consisting of hydroxy, halogen, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, -O-phenyl, linear or branched C₁-C₆ perfluoroalkyl, linear or branched C₁-C₆ perfluoroalkoxy, an optionally at least mono-substituted phenyl and 5- to 6-membered heteroaryl, more preferably from the group consisting of halogen, linear or branched C₁-C₆ alkyl, -O-phenyl, optionally at least mono-substituted phenyl and 5- to 6-membered heteroaryl, even more preferably from the group consisting of fluorine, chlorine, -O-phenyl, linear or branched C₁-C₆ alkyl, optionally at least mono-substituted phenyl and 5- to 6-membered heteroaryl.

If one or more of the rings of a mono- or polycyclic aromatic ring system contain one or more heteroatoms, these heteroatoms – like the heteroatoms of the previously mentioned 5- to 6-membered heteroaryl – may preferably be chosen from the group consisting of nitrogen, sulphur and oxygen.

5

If the previously mentioned phenyl radical is itself substituted by one or more substituents, each one of these substituents may preferably be chosen from the group consisting of fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, trifluoromethyl radical, cyano radical and a -NR¹²R¹³ radical, wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁-C₆ alkyl.

10

If the previously mentioned alkylene, alkenylene or alkynylene group is substituted by one or more substituents, each of these substituents may preferably be chosen from the group consisting of hydroxy, halogen, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ perfluoroalkyl, linear or branched C₁-C₆ perfluoroalkoxy or an optionally at least mono-substituted phenyl radical. If said phenyl radical is itself substituted by one or more substituents, each one of these substituents may preferably be chosen from the group consisting of fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, trifluoromethyl radical, cyano radical and a -NR¹²R¹³ radical, wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁-C₆ alkyl.

15

20

25

If one or more of the substituents R², R³, R⁴, R⁵ and R⁷ represents an alkoxy radical, said radical may have 1 to 6, preferably 1 to 3 carbon atoms.


30

Those skilled in the art understand that the term “condensed” indicates that the condensed rings share more than one atom. The terms “annulated” or “fused” may also be used for this type of bonding.

-12-

Sulfonamide derivatives of general formula (Ia) are preferred, wherein R^1 represents a $-NR^8R^9$ radical or a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing 5- or 6-membered cycloaliphatic radical, which may be condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system, whereby the rings of the ring system are 5- or 6-membered,

more preferably R^1 represents an $-NR^8R^9$ radical or a radical chosen from the group consisting of



wherein, if present, the dotted line represents an optional chemical bond, and R^{10} represents hydrogen, a linear or branched C_1 - C_6 alkyl radical or a benzyl radical, preferably hydrogen or a C_1 - C_2 alkyl radical and R^2 to R^9 , A and n are defined as above.

Also preferred are sulfonamide derivatives of general formula (Ia), wherein R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen, a linear or branched C_1 - C_6 alkoxy radical, a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_6 alkenyl radical, or a linear or branched, optionally at least mono-substituted C_2 - C_6 alkynyl radical,

more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen or a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical,

5

even more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen,

and R^1 , R^6 , R^8 , R^9 , A and n are defined as above.

10

Sulfonamide derivatives of general formula (Ia) are also preferred, wherein R^6 , represents hydrogen, a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_6 alkenyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_6 alkynyl radical,

15

more preferably R^6 represents hydrogen or a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical,

20

even more preferably R^6 represents hydrogen or a C_1 - C_2 alkyl radical,

and R^1 - R^5 , R^7 - R^9 , A and n are defined as above.

25

Furthermore, sulfonamide derivatives of general formula (Ia) are also preferred, wherein R^8 and R^9 , identical or different, each represent hydrogen, a linear or branched, optionally at least mono-substituted C_1 - C_{10} alkyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_{10} alkenyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_{10} alkynyl radical,

30

with the proviso that R^8 and R^9 do not represent hydrogen at the same time, and if one of them, R^8 and R^9 , represents a saturated or unsaturated, linear or branched, optionally at least mono-substituted C_1 - C_4 aliphatic radical,

the other one represents a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical with at least five carbon atoms,

5 or

R^8 and R^9 , together with the bridging nitrogen form a saturated or unsaturated, optionally at least mono-substituted, optionally at least one further heteroatom as a ring member containing 5- or 6-membered heterocyclic ring, which may be
10 condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system, whereby the rings of the ring system are 5- 6- or 7-membered,
and R^1 - R^7 , A and n are defined as above.

15


Particularly preferred are sulfonamide derivatives of general formula (Ia), wherein R^8 and R^9 , identical or different, each represent hydrogen or a linear or branched C_1 - C_{10} alkyl radical,

20 with the proviso that R^8 and R^9 are not hydrogen at the same time, and if one of them, R^8 or R^9 , is a saturated or unsaturated, linear or branched, optionally at least mono-substituted C_1 - C_4 aliphatic radical, the other one of them is a saturated or unsaturated, linear or branched, optionally at least mono-substituted aliphatic radical with at least five carbon atoms,

25

or

R^8 and R^9 together with the bridging nitrogen atom form a radical chosen from the group consisting of



wherein R^{11} represents hydrogen, a linear or branched C_1 - C_6 alkyl radical or a benzyl radical, preferably hydrogen or a C_1 - C_2 alkyl radical, and R^1 - R^7 , A and n are defined as above.

5


Furthermore, sulfonamide derivatives of general formula (Ia) are preferred, wherein A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered, which may be bonded via an optionally at least mono-substituted C_1 - C_6 alkylene group, an optionally at least mono-substituted C_2 - C_6 alkenylene group or an optionally at least mono-substituted C_2 - C_6 alkynylene group and/or wherein the ring(s) may contain at least one heteroatom as a ring member,

preferably A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered and wherein one or more of the rings contain at least one heteroatom,

15


or a radical chosen from the group consisting of

-16-



- wherein X, Y, Z, independently from one another, each represent a radical
 5 selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C_1 - C_6 alkyl, linear or branched C_1 - C_6 alkoxy, linear or branched C_1 - C_6 alkylthio, a trifluoromethyl radical, a cyano radical and a $-NR^{12}R^{13}$ radical,
- 10 wherein R^{12} and R^{13} , identical or different, each represent hydrogen or linear or branched C_1 - C_6 alkyl,
- W represents a single chemical bond between the two rings, a CH_2 , O, S group or a NR^{14} radical,
- 15 wherein R^{14} is hydrogen or a linear or branched C_1 - C_6 alkyl,
- m is 0, 1, 2, 3 or 4 and
- 20 m1 is 1 or 2, preferably 2, and R^1 - R^9 and n are defined as above.

- Furthermore, sulfonamide derivatives of general formula (Ia) are preferred, wherein A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered, which may be bonded via an optionally at least mono-substituted C₁–C₆ alkylene group, an optionally at least mono-substituted C₂–C₆ alkenylene group or an optionally at least mono-substituted C₂–C₆ alkynylene group, and/or wherein the ring(s) may contain at least one heteroatom as a ring member,
- more preferably A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered and wherein one or more of the rings contain at least one heteroatom, or a radical chosen from the group consisting of



- wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C₁–C₆ alkyl, linear or branched C₁–C₆ alkoxy, linear or branched C₁–C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -NR¹²R¹³ radical,
- wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁–C₆ alkyl,

W represents a single chemical bond between the two rings, a CH₂, O, S group

-18-

or a NR^{14} radical,

wherein R^{14} is hydrogen or a linear or branched $\text{C}_1\text{-C}_6$ alkyl,

5 and

m is 0, 1, 2, 3 or 4.


and $\text{R}^1\text{-R}^9$ and n are defined as above.

10

Furthermore, sulfonamide derivatives of general formula (Ia) are preferred, wherein A represents an heteroaryl radical selected from the group consisting of benzo[b]thiophenyl and imidazo[2,1-b]thiazolyl which may be substituted by 1, 2 or 3 substituents selected from the group consisting of chlorine, methyl, phenyl and -O-phenyl and/or which may be bonded via a C_{1-2} alkylene group,

15

or a radical chosen from the group consisting of



20

-19-

- wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -
- 5 NR¹²R¹³ radical,
- wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁-C₆ alkyl,
- 10 W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,
- wherein R¹⁴ is hydrogen or a linear or branched C₁-C₆ alkyl,
- 15 m is 0, 1, 2, 3 or 4 and
- m₁ is 1 or 2, preferably 2, and R¹-R⁹ and n are defined as above.
- Furthermore sulfonamide derivatives of general formula (Ia) are preferred,
- 20 wherein n is 0, 1, 2, 3 or 4; preferably n is 1 or 2; more preferably n is 2 and R¹ to R⁹ and A are defined as above.
- Those most preferred compounds of general formula (Ia) are selected
- 25 from the group consisting of
- [9] 5-Chloro-3-methyl-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-benzo[b]thiophene-2-sulfonamide,
- 30 [10] N-(1-[2-(Pyrrolidin-1-yl)ethyl]-1H-indol-6-yl)-naphthalene-2-sulfonamide,
- [11] N-[1-[2-Pyrrolidin-1-yl]ethyl]-1H-indol-6-yl]-naphthalene-1-sulfonamide,

[12] 6-Chloro-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-imidazo[2,1-b]thiazole-5-sulfonamide,

5 [13] 4-Phenyl-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-benzenesulfonamide

[14] 2-(Naphth-1-yl)-N-(1-(2-(pyrrolidin-1-yl)-ethyl)-1H-indol-6-yl)-ethansulfonamide,

10

[15] 4-Phenoxy-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-benzenesulfonamide and

[16] 3,5-Dichloro-N-(1-(2-(pyrrolidin-1-yl)-1H-indol-6-yl)-benzenesulfonamide,

15

and their corresponding salts and solvates.

Furthermore, sulfonamide derivatives of general formula (Ib) are also preferred, wherein R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each
20 represent hydrogen, C_1 - C_6 alkoxy radical, a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_6 alkenyl radical, or a linear or branched, optionally at least mono-substituted C_2 - C_6 alkynyl radical,

25 more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen or a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical,

even more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each
30 represent hydrogen,

and R^1 , R^6 , R^8 , R^9 , A and n are defined as above.

Sulfonamide derivatives of general formula (Ib) are also preferred, wherein R^6 represents hydrogen, a linear or branched, optionally at least mono-substituted C_1-C_6 alkyl radical, a linear or branched, optionally at least mono-substituted C_2-C_6 alkenyl radical, a linear or branched, optionally at least mono-substituted C_2-C_6 alkynyl radical,

more preferably R^6 represents hydrogen or a linear or branched, optionally at least mono-substituted C_1-C_6 alkyl radical,

10

more preferably R^6 represents hydrogen or a C_1-C_2 alkyl radical,

and R^1-R^5 , R^7-R^9 , A and n are defined as above.

Furthermore, sulfonamide derivatives of general formula (Ib) are also preferred, wherein R^8 and R^9 , identical or different, each represent hydrogen or a linear or branched, optionally at least mono-substituted C_1-C_4 alkyl radical, and R^1-R^7 , A and n are defined as above.

Particularly preferred are sulfonamide derivatives of general formula (Ib) wherein R^8 and R^9 , identical or different, each represent hydrogen or a C_1-C_2 alkyl radical, with the proviso that R^8 and R^9 are not hydrogen at the same time, and R^1-R^7 , A and n are defined as above.


Furthermore, sulfonamide derivatives of general formula (Ib) are preferred, wherein A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered, which may be bonded via an optionally at least mono-substituted C_1-C_6 alkylene group, an optionally at least mono-substituted C_2-C_6 alkenylene group or an optionally at least mono-substituted C_2-C_6 alkynylene group and/or wherein the ring(s) may contain at least one heteroatom as a ring member,

30

-22-

preferably A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered and wherein one or more of the rings contain at least one heteroatom,

- 5 or a radical chosen from the group consisting of



- wherein X, Y, Z, independently from one another, each represent a radical
 10 selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -NR¹²R¹³ radical,

- 15 wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁-C₆ alkyl,


W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,

wherein R^{14} is hydrogen or a linear or branched C_1 - C_6 alkyl,

m is 0, 1, 2, 3 or 4 and

- 5 m1 is 1 or 2, preferably 2 and R^1 - R^9 and n are defined as above.

- Furthermore, sulfonamide derivatives of general formula (Ib) are preferred, wherein A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered,
 10 which may be bonded via an optionally at least mono-substituted C_1 - C_6 alkylene group, an optionally at least mono-substituted C_2 - C_6 alkenylene group or an optionally at least mono-substituted C_2 - C_6 alkynylene group, and/or wherein the ring(s) may contain at least one heteroatom as a ring member,
 15 more preferably A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered and wherein one or more of the rings contain at least one heteroatom, or a radical chosen from the group consisting of



- 20 wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C_1 - C_6 alkyl, linear or branched C_1 - C_6 alkoxy, linear or branched C_1 - C_6 alkylthio, a trifluoromethyl radical, a cyano radical and a - $NR^{12}R^{13}$ radical,

wherein R^{12} and R^{13} , identical or different, each represent hydrogen or linear or branched C_1 - C_6 alkyl,

- 5 W represents a single chemical bond between the two rings, a CH_2 , O, S group or a NR^{14} radical,

wherein R^{14} is hydrogen or a linear or branched C_1 - C_6 alkyl,

- 10 and

m is 0, 1, 2, 3 or 4.


and R^1 - R^9 and n are defined as above.

15


Furthermore, sulfonamide derivatives of general formula (lb) are preferred, wherein A represents an heteroaryl radical selected from the group consisting of benzo[b]thiophenyl and imidazo[2,1-b]thiazolyl which may be substituted by 1, 2 or 3 substituents selected from the group consisting of chlorine, methyl, phenyl

- 20 and -O-phenyl and/or which may be bonded via a C_{1-2} alkylene group,

or a radical chosen from the group consisting of



-25-



wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine,
 5 linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -NR¹²R¹³ radical,

wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or
 10 branched C₁-C₆ alkyl,

W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,

15 wherein R¹⁴ is hydrogen or a linear or branched C₁-C₆ alkyl,

m is 0, 1, 2, 3 or 4 and

m₁ is 1 or 2, preferably 2 and R¹-R⁹ and n are defined as above
 20 Furthermore sulfonamide derivatives of general formula (Ib) are preferred, wherein n is 0, 1, 2, 3 or 4; preferably n is 1 or 2; more preferably n is 2 and R¹ to R⁹ and A are defined as above.

Those most preferred compounds are sulfonamide derivatives of general
 25 formula (Ib), selected from the group consisting of

[1] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-5-chloro-3-methylbenzo[b]thiophene-2-sulfonamide,

- [2] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-naphthalene-2-sulfonamide,
- [3] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-naphthalene-1-sulfonamide,
- 5 [4] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-6-chloroimidazo[2,1-b]thiazole-5-sulfonamide,
- [5] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-4-phenylbenzenesulfonamide,
- 10 [6] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-2-(naphthalene-1-yl)-ethanesulfonamide,
- [7] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-4-phenoxybenzenesulfonamide,
- 15 [8] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-3,5-dichlorobenzenesulfonamide,

and their corresponding salts and solvates.


20

Furthermore, sulfonamide derivatives of general formula (Ic) are preferred, wherein R^1 represents a $-NR^8R^9$ radical or a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing 5- or 6-membered cycloaliphatic radical, which may be

25 condensed with a saturated or unsaturated, optionally at least mono-substituted, optionally at least one heteroatom as a ring member containing mono- or bicyclic cycloaliphatic ring system, whereby the rings of the ring system are 5- or 6-membered,

30 more preferably R^1 represents an $-NR^8R^9$ radical or a radical chosen from the group consisting of

-27-



wherein, if present, the dotted line represents an optional chemical bond, and R^{10} represents hydrogen, a linear or branched C_1 - C_6 alkyl radical or a benzyl radical, preferably hydrogen or a C_1 - C_2 alkyl radical and R^2 to R^9 , A and n are defined as above.

Also preferred are sulfonamide derivatives of general formula (Ic), wherein R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen, a linear or branched C_1 - C_6 alkoxy radical, a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical, a linear or branched, optionally at least mono-substituted C_2 - C_6 alkenyl radical, or a linear or branched, optionally at least mono-substituted C_2 - C_6 alkynyl radical,

more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen or a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical,

even more preferably R^2 , R^3 , R^4 , R^5 and R^7 , identical or different, each represent hydrogen,

and R^1 , R^6 , R^8 , R^9 , A and n are defined as above.

-28-

Sulfonamide derivatives of general formula (Ic) are also preferred,
wherein R^6 , represents hydrogen, a linear or branched, optionally at least
mono-substituted C_1 - C_6 alkyl radical, a linear or branched, optionally at least
mono-substituted C_2 - C_6 alkenyl radical, a linear or branched, optionally at least
5 mono-substituted C_2 - C_6 alkynyl radical,

more preferably R^6 represents hydrogen or a linear or branched, optionally at
least mono-substituted C_1 - C_6 alkyl radical,
10 even more preferably R^6 represents hydrogen or a C_1 - C_2 alkyl radical,
and R^1 - R^5 , R^7 - R^9 , A and n are defined as above.

Furthermore, sulfonamide derivatives of general formula (Ic) are also preferred,
15 wherein R^8 and R^9 , identical or different, each represent hydrogen, a linear or
branched, optionally at least mono-substituted C_1 - C_{10} alkyl radical, a linear or
branched, optionally at least mono-substituted C_2 - C_{10} alkenyl radical, a linear
or branched, optionally at least mono-substituted C_2 - C_{10} alkynyl radical,

20 or


R^8 and R^9 , together with the bridging nitrogen form a saturated or unsaturated,
optionally at least mono-substituted, optionally at least one further heteroatom
as a ring member containing 5- or 6-membered heterocyclic ring, which may be
25 condensed with a saturated or unsaturated, optionally at least mono-
substituted, optionally at least one heteroatom as a ring member containing
mono- or bicyclic cycloaliphatic ring system, whereby the rings of the ring
system are 5- 6- or 7-membered,
and R^1 - R^7 , A and n are defined as above.

30

Particularly preferred are sulfonamide derivatives of general formula (Ic), wherein R^8 and R^9 , identical or different, each represent hydrogen or a linear or branched C_1 - C_{10} alkyl radical,

or

- 5 R^8 and R^9 together with the bridging nitrogen atom form a radical chosen from the group consisting of




- wherein R^{11} represents hydrogen, a linear or branched C_1 - C_6 alkyl radical or a benzyl radical, preferably hydrogen or a C_1 - C_2 alkyl radical,
 10 and R^1 - R^7 , A and n are defined as above.

- Furthermore, sulfonamide derivatives of general formula (Ic) are preferred, wherein A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered,
 15 which may be bonded via an optionally at least mono-substituted C_1 - C_6 alkylene group, an optionally at least mono-substituted C_2 - C_6 alkenylene group or an optionally at least mono-substituted C_2 - C_6 alkynylene group and/or wherein the ring(s) may contain at least one heteroatom as a ring member,
 20 preferably A represents an optionally at least mono-substituted mono- or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered and wherein one or more of the rings contain at least one heteroatom,

-30-

or a radical chosen from the group consisting of



- 5 wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine, linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -NR¹²R¹³ radical,

10

wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or branched C₁-C₆ alkyl,

- 15 W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,


wherein R¹⁴ is hydrogen or a linear or branched C₁-C₆ alkyl,

- 20 m is 0, 1, 2, 3 or 4 and

m1 is 1 or 2, preferably 2 and R^1 - R^9 and n are defined as above.

Furthermore, sulfonamide derivatives of general formula (Ic) are
 5 preferred, wherein A represents an optionally at least mono-substituted mono-
 or polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered,
 which may be bonded via an optionally at least mono-substituted C_1 - C_6
 alkylene group, an optionally at least mono-substituted C_2 - C_6 alkenylene group
 or an optionally at least mono-substituted C_2 - C_6 alkynylene group, and/or
 10 wherein the ring(s) may contain at least one heteroatom as a ring member,

more preferably A represents an optionally at least mono-substituted mono- or
 polycyclic aromatic ring system, wherein the ring(s) is/are 5- or 6-membered
 and wherein one or more of the rings contain at least one heteroatom,
 15 or a radical chosen from the group consisting of



wherein X, Y, Z, independently from one another, each represent a radical
 selected from the group consisting of hydrogen, fluorine, chlorine, bromine,
 linear or branched C_1 - C_6 alkyl, linear or branched C_1 - C_6 alkoxy, linear or
 20 branched C_1 - C_6 alkylthio, a trifluoromethyl radical, a cyano radical and a -
 $NR^{12}R^{13}$ radical,

wherein R^{12} and R^{13} , identical or different, each represent hydrogen or linear or
 branched C_1 - C_6 alkyl,

W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,

5 wherein R¹⁴ is hydrogen or a linear or branched C₁-C₆ alkyl,

and


m is 0, 1, 2, 3 or 4.

10


and R¹-R⁹ and n are defined as above.

Furthermore, sulfonamide derivatives of general formula (Ic) are preferred, wherein A represents an heteroaryl radical selected from the group consisting of
 15 benzo[b]thiophenyl and imidazo[2,1-b]thiazolyl which may be substituted by 1, 2 or 3 substituents selected from the group consisting of chlorine, methyl, phenyl and -O-phenyl and/or which may be bonded via a C₁₋₂ alkylene group,

or a radical chosen from the group consisting of



-33-



wherein X, Y, Z, independently from one another, each represent a radical selected from the group consisting of hydrogen, fluorine, chlorine, bromine,
 5 linear or branched C₁-C₆ alkyl, linear or branched C₁-C₆ alkoxy, linear or branched C₁-C₆ alkylthio, a trifluoromethyl radical, a cyano radical and a -NR¹²R¹³ radical,

wherein R¹² and R¹³, identical or different, each represent hydrogen or linear or
 10 branched C₁-C₆ alkyl,


W represents a single chemical bond between the two rings, a CH₂, O, S group or a NR¹⁴ radical,

15 wherein R¹⁴ is hydrogen or a linear or branched C₁-C₆ alkyl,

m is 0, 1, 2, 3 or 4 and

m₁ is 1 or 2, preferably 2 and R¹-R⁹ and n are defined as above
 20 Furthermore sulfonamide derivatives of general formula (lc) are preferred, wherein n is 0, 1, 2, 3 or 4; preferably n is 1 or 2; more preferably n is 2 and R¹ to R⁹ and A are defined as above.

Another aspect of the present invention are compounds of general formula (Ic),



(Ic),

5

wherein

R¹ represents a -NR⁸R⁹ radical,

10 R², R³, R⁴, R⁵ and R⁷ each represent hydrogen,

R⁶ represents hydrogen,

15 R⁸ and R⁹, identical or different, each represent methyl, ethyl, n-propyl or n-propyl, more preferably methyl,

or

20 R⁸ and R⁹, together with the bridging nitrogen atom form a 5- or 6-membered heterocyclic ring, more preferably form a pyrrolidine ring or a piperidine ring

and

25 A represents an aryl or heteroaryl radical selected from the group consisting of phenyl, naphthyl, benzo[b]thiophenyl and imidazo[2,1-b]thiazolyl which may be substituted by 1, 2 or 3 substituents selected from the group consisting of chlorine, methyl, phenyl and -O-phenyl and/or which may be bonded via a C₁₋₂

alkylene group,

and n is 2,

5 optionally in form of one of its stereoisomers, preferably enantiomers or diastereomers, a racemate or in form of a mixture of at least two of its stereoisomers, preferably enantiomers and/or diastereomers, in any mixing ratio, or a salt thereof, preferably a corresponding, physiologically acceptable salt thereof, or a corresponding solvate thereof.

10

Those most preferred compounds are sulfonamide derivatives of general formula (Ic), selected from the group consisting of

- 15 [1] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-5-chloro-3-methylbenzo[b]thiophene-2-sulfonamide,
- [2] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-naphthalene-2-sulfonamide,
- [3] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-naphthalene-1-sulfonamide,
- 20 [4] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-6-chloroimidazo[2,1-b]thiazole-5-sulfonamide,
- [5] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-4-phenylbenzenesulfonamide,
- 25 [6] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-2-(naphthalene-1-yl)-ethanesulfonamide,
- [7] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-4-
- 30 phenoxybenzenesulfonamide,


- [8] N-[1-(2-Dimethylaminoethyl)-1H-indol-6-yl]-3,5-dichlorobenzenesulfonamide,
- [9] 5-Chloro-3-methyl-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-benzo[b]thiophene-2-sulfonamide,
- [10] N-(1-[2-(Pyrrolidin-1-yl)ethyl]-1H-indol-6-yl)-naphthyl-2-sulfonamide,
- [11] N-[1-[2-Pyrrolidin-1-yl]ethyl]-1H-indol-6-yl]-naphthalene-1-sulfonamide,
- [12] 6-Chloro-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-imidazo[2,1-b]thiazole-5-sulfonamide,
- [13] 4-Phenyl-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-benzenesulfonamide
- [14] 2-(Naphth-1-yl)-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-ethansulfonamide,
- [15] 4-Phenoxy-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-benzenesulfonamide and
- [16] 3,5-Dichloro-N-(1-(2-(pyrrolidin-1-yl)-1H-indol-6-yl)-benzenesulfonamide,
- and their corresponding salts and solvates.

The present invention likewise refers to the salts, preferably the physiologically acceptable salts of the compounds of general formula (Ia) and/or (Ib) and/or of general formula (Ic), preferably the addition salts of mineral acids, more preferably of hydrochloric acid, hydrobromic acid, phosphoric acid, sulphuric acid, nitric acid, and the salts of organic acids, more preferably of citric acid, maleic acid, fumaric acid, tartaric acid or their derivatives, *p*-

toluenesulphonic acid, methanesulphonic acid, camphorsulphonic acid, etc.


Below, the expression sulfonamide derivatives of general formula (I) refers to one or more compounds of general formula (Ia) and/or to one or more
 5 compounds of general formula (Ib) and/or to one or more compounds of general formula (Ic), respectively, and optionally in form of one of their stereoisomers, preferably enantiomers or diastereomers, their racemate, or in form of a mixture of at least two of their stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a salt thereof, preferably a corresponding physiologically
 10 acceptable salt thereof, or a corresponding solvate thereof.

Another aspect of the present invention consists of a process for preparing the new derivatives of general formula (I), wherein R^1 - R^9 , n and A have the previously indicated meaning, according to which at least one
 15 compound of general formula (II),



(II)

wherein A has the previously mentioned meaning, and X is an acceptable leaving group, preferably an halogen atom, more preferably chlorine; is reacted with at least one substituted 6-aminoindole of general formula (III), or one of its
 20 suitably protected derivatives;



(III)

wherein R^1 - R^7 and n have the previously indicated meaning, and, if necessary, the protective groups are removed in order to obtain the corresponding

sulfonamide derivative of formula (I), which can be purified and/or isolated by means of conventional methods known in the prior art.

The reaction between the compounds of general formula (II) and (III) is usually carried out in the presence of an organic reaction medium, preferably in the presence of dialkyl ether, more preferably diethyl ether or a cyclic ether, more preferably tetrahydrofuran or dioxane, an halogenated organic hydrocarbon, more preferably methylene chloride or chloroform, an alcohol, more preferably methanol or ethanol, a dipolar aprotic solvent, more preferably acetonitrile, pyridine or dimethylformamide, or any other suitable reaction medium. Naturally, mixtures of at least two of the classes of the mentioned compounds or at least two compounds of one class may also be used.


The reaction is preferably carried out in the presence of a suitable base, for example, an inorganic base, more preferably alkaline metal hydroxides and alkaline metal carbonates, or in the presence of an organic base, more preferably triethylamine, N-ethyldiisopropylamine or pyridine.

The most suitable reaction temperatures range from 0 °C to room temperature, that is, approximately 25 °C, and the reaction time preferably comprises from 5 minutes to 24 hours.

The resulting sulfonamide derivative of general formula (I) can be purified and/or isolated according to conventional methods known in the prior art.

Preferably, the sulfonamide derivatives of general formula (I) can be isolated by evaporating the reaction medium, adding water and, if necessary, adjusting the pH so that a solid which can be isolated by filtration is obtained; or it can be extracted with a water immiscible solvent, preferably chloroform, and be purified by chromatography or recrystallization of a suitable solvent.

The compounds of general formula (II) are commercially available, or they can be prepared according to standard methods known in the prior art, for example by methods similar to those described in the literature [E.E.Gilbert, Synthesis, 1969, 1, 3]. The compounds of general formula (III) can also be prepared according to standard methods known in the prior art, for example by methods similar to those described in the literature [Ham, Peter; Gaster, Laramie Mary; King, Francis David; Duckworth, David Malcolm. Preparation of N-heteroaryl-4'-oxadiazolylbiphenylcarboxamides as 5HT1D antagonists. WO 9532967 A1 19951207; Basanagoudar, L.D.; Siddappa, S. Cyanoethylation of nitroindoles. Journal of the Indian Chemical Society (1972), 49 (8), 811-13.; Chen, Guoqing; Adams, Jeffrey; Bemis, Jean; Booker, Shon; Cai, Guolin; Croghan, Michael; Dipietro, Lucian; Dominguez, Celia; Elbaum, Daniel; Germain, Julie; Geuns-Meyer, Stephanie; Handley, Michael; Huang, Qi; Kim, Joseph L.; Kim, Tae-seong; Kiselyov, Alexander; Ouyang, Xiaohu; Patel, Vinod F.; Smith, Leon M.; Stec, Markian; Tasker, Andrew; Xi, Ning; Xu, Shimin; Yuan, Chester Chenguang. Preparation of heterocyclalkylamine derivatives as remedies for angiogenesis mediated diseases. WO 0266470 A1 20020829. European Journal of Medicinal Chemistry, 23 (4), 373-7; 1988]. One of them consists of nitro group reduction of derivatives of general formula (IV) by methods known in the art, as for example YAMASHKIN, S.A.; YUROVSKAYA, M.A.; Chem Heterocycl Compd (N.Y.), 1999, 35 (12), 1426-1432. BOOTHROYD, S.R.; KERR, M.A.; Tetrahedron Lett, 1995, 36 (14), 2411-2414. MACOR, J.E.; POST, R.; RYAN, K.; Synth Common, 1993, 23 (1), 65-72,



(IV)


wherein R¹-R⁷ and n have the previously indicated meaning, or one of their suitably protected derivatives, and, if necessary, the protective groups are

-40-

removed in order to obtain the corresponding amine of general formula (III), which can be purified and/or isolated by means of conventional methods known in the prior art.

- 5 The compounds of general formula (IV) can also be prepared according to standard methods known in the prior art, for example by methods similar to those described in the European Journal of Medicinal Chemistry, 23 (4), 373-7; 1988; Farmaco, 51 (1), 75-8; 1996; Heterocycles, 55 (6), 1151-1159; 2001; Ham, Peter; Gaster, Laramie Mary; King, Francis David; Duckworth, David
- 10 Malcolm. Preparation of N-heteroaryl-4'-oxadiazolylbiphenylcarboxamides as 5HT1D antagonists, WO 9532967 A1 19951207.

- One of them consists in the alkylation of nitro derivatives of general formula (IV) by methods known in the art: MACCHIA, M.; MANERA, C.;
- 15 NENCETTI, S.; ROSSELLO, A.; BROCCALI, G.; LIMONTA, D.; Farmaco, Ed Sci [FRPSAX] 1996, 51 (1), 75-78. BHAGWAT, S. S.; GUDE, C.; Tetrahedron Lett, 1994, 35 (12), 1847-1850. BRATTON, L.D.; ROTH, B.D.; TRIVEDI, B.K.; UNANGST, P.C.; J Heterocycl Chem, 2000, 37 (5), 1103-1108,



(V)

- 20 wherein R²-R⁷ and n have the previously mentioned meaning, or one of their suitably protected derivatives, and, if necessary, the protective groups are removed in order to obtain the corresponding amine of general formula (III), which can be purified and/or isolated by means of conventional methods known in the prior art.

25

The compounds of general formula (V) are commercially available or can also be prepared according to standard methods known in the prior art, as for example OTTONI, O.; CRUZ, R.; KRAMMER, N.H.; Tetrahedron Lett [TELEAY] 1999, 40 (6), 1117-1120. VOROB'EVA, S.L.; BUYANOV, V.N.; SUVOROV, N.N.; Khim Geterosikl Soedin [KGSSAQ] 1991, (5), 636-637. KATRITZKY, A.R.; RACHWAL, S.; BAYYUK, S.; Org Prep Proceed Int [OPPIAK] 1991, 23 (3), 357-363. MOSKALEV, N.; MAKOSZA, M.; Heterocycles [HTCYAM] 2000, 52 (2), 533-536.

10 The respective literature descriptions are incorporated by reference and form part of the disclosure.

Another aspect of the present invention consists in a process for preparing the new sulfonamide derivatives of general formula (I), wherein R^1 - R^5 , R^7 - R^9 , A and n have the previously indicated meaning and R^6 is an alkyl radical, preferably a linear or branched, optionally at least mono-substituted C_1 - C_6 alkyl radical, by alkylation of a sulfonamide derivative of general formula (I), wherein R^1 - R^5 , R^7 - R^9 , n and A have the previously indicated meaning, and R^6 is an hydrogen atom, with an alkyl halogenide or dialkyl sulfate.

20

The alkylation reaction is carried out preferably in the presence of a suitable base, more preferably in the presence of alkaline metal hydroxides and alkaline metal carbonates, metal hydrides, metal alkoxides, even more preferably sodium methoxide or potassium tert-butoxide, organometallic compounds, even more preferably butyllithium or tert-butyllithium, in the presence of an organic reaction medium, more preferably dialkyl ether, even more preferably diethyl ether, or a cyclic ether, even more preferably tetrahydrofuran or dioxane, an hydrocarbon, even more preferably toluene, an alcohol, even more preferably methanol or ethanol, a dipolar aprotic solvent, even more preferably acetonitrile, pyridine or dimethylformamide, or any other suitable reaction medium. Naturally, mixtures of at least two of the classes of the mentioned compounds or at least two compounds of one class may also be

30

used.

The most suitable reaction temperatures range from 0°C to the boiling temperature of the reaction medium, and the reaction times are preferably
5 comprised from 1 to 24 hours.

Preferably, the resulting sulfonamide derivative of general formula (I) can be isolated by filtration, concentrating the filtrate under reduced pressure, adding water and, if necessary, adjusting the pH so that a solid which can be
10 isolated by filtration is obtained; or it can be extracted with a water immiscible solvent, preferably chloroform, and be purified by chromatography or recrystallization of a suitable solvent.

The salts, preferably pharmaceutically acceptable salts of the
15 compounds of general formula (I), may be prepared by means of conventional methods known in the prior art, preferably by reaction with a mineral acid, more preferably by reaction with hydrochloric acid, hydrobromic acid, phosphoric acid, sulphuric acid or nitric acid, or by reaction with organic acids, more preferably by reaction with citric acid, maleic acid, fumaric acid, tartaric
20 acid, or their derivatives, *p*-toluenesulphonic acid, methanesulphonic acid, camphorsulphonic acid, etc., in a suitable solvent, preferably methanol, ethanol, diethyl ether, ethyl acetate, acetonitrile or acetone, and obtaining the resulting salts by using the usual techniques for the precipitation or crystallization of the corresponding salts.

25

The preferred physiologically acceptable salts of the sulfonamide derivatives of general formula (I) are the addition salts of mineral acids, more preferably of hydrochloric acid, hydrobromic acid, phosphoric acid, sulphuric acid or nitric acid, and the addition salts of organic acids, more preferably
30 citric acid, maleic acid, fumaric acid, tartaric acid, or their derivatives, *p*-toluenesulphonic acid, methanesulphonic acid, camphorsulphonic acid, etc.

The solvates, preferably the physiologically acceptable solvates, more preferably hydrates, of the sulfonamide derivatives of general formula (I) or of the corresponding physiologically acceptable salts, may be prepared by methods known in the prior art.

5

During some of the synthetic sequences described or in the preparation of the suitable reagents used, it may be necessary and/or desirable to protect sensitive or reactive groups in some of the molecules used. This can be done by means of using conventional protective groups preferably those described in the literature [Protective groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Chemistry, John Wiley & Sons, 1991]. The protective groups can be removed in a suitable subsequent stage by methods known in the prior art. The respective literature descriptions are incorporated by reference and form part of the disclosure.

10
15

If the sulfonamide derivatives of general formula (I) are obtained in the form of a mixture of stereoisomers, preferably enantiomers or diastereomers, said mixtures can be separated by means of standard processes known in the prior art, for example chromatographic methods or crystallization with chiral agents.

20

Another aspect of the present invention is a medicament comprising at least one indol-6-yl sulfonamide derivative of general formula (I), optionally in the form of one of its stereoisomers, preferably enantiomer or diastereomer, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, and optionally at least one or more pharmaceutically acceptable adjuvants.

25
30

This medicament is suitable for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, 5 cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar 10 disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder), and other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in 15 animals, preferably in mammals, more preferably in humans.

Another aspect of the present invention is a medicament comprising at least one indol-6-yl sulfonamide derivative of general formula (Ia), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, 20 its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, and optionally at least one or more pharmaceutically acceptable 25 adjuvants.

This medicament is suitable for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, 30 cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome,

for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder),

more suitable for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome.

15

Another aspect of the present invention is a medicament comprising at least one indol-6-yl sulfonamide derivative of general formula (Ib), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, and optionally at least one or more pharmaceutically acceptable adjuvants.

20

This medicament is suitable for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of

30

the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis
5 or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder),

more suitable for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes,
10 neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) in humans and/or in animals, preferably in mammals, more preferably in humans.

15 Another aspect of the present invention is a medicament comprising at least one indol-6-yl sulfonamide derivative of general formula (Ic), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a
20 corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, and optionally at least one or more pharmaceutically acceptable adjuvants.

This medicament is suitable for 5-HT₆ receptor regulation, for the prophylaxis
25 and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or
30 treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar

disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder), in
5 humans and/or in animals, preferably in mammals, more preferably in humans.

The medicament obtained according to the present invention is particularly suitable for the administration to mammals, including humans. The medicament can preferably be administered to all ages, namely children,
10 adolescents and adults.

Another aspect of the present invention is the use of at least one sulfonamide derivative of general formula (I), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in
15 form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ration, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof for the manufacture of a medicament for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the
20 regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome,
25 for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder), and
30 other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in animals, preferably in mammals, more preferably in humans.

Another aspect of the present invention is the use of at least one sulfonamide derivative of the previous general formula (Ia), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or
5 diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, for the manufacture of a medicament for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body
10 weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of
15 the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) in
20 humans and/or in animals, preferably in mammals, more preferably in humans,

preferably for the manufacture of a medicament for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance,
25 increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome.

Another aspect of the present invention is the use of at least one sulfonamide derivative of the previous general formula (Ib), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably
5 enantiomers or diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, for the manufacture of a medicament for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or
10 reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of
15 disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder), and
20 other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in animals, preferably in mammals, more preferably in humans,

preferably for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders,
25 depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) and other disorders mediated by the 5-HT₆ serotonin
30 receptor in humans and/or in animals, preferably in mammals, more preferably in humans.

Another aspect of the present invention is the use of at least one sulfonamide derivative of the previous general formula (Ic), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a corresponding physiologically acceptable salt thereof or a corresponding solvate thereof, for the manufacture of a medicament for 5-HT₆ receptor regulation, for the prophylaxis and/or treatment of a disorder or disease related to food intake, preferably for the regulation of appetite, for the maintenance, increase or reduction of body weight, for the prophylaxis and/or treatment of obesity, bulimia, anorexia, cachexia or type II diabetes (non insulin dependent diabetes mellitus), preferably type II diabetes caused by obesity, for the prophylaxis and/or treatment of gastrointestinal tract disorders, preferably irritable bowel syndrome, for cognitive enhancement, for the prophylaxis and/or treatment of disorders of the central nervous system, anxiety, panic disorders, depression, bipolar disorders, cognitive memory disorders, senile dementia processes, neurodegenerative disorders, preferably Alzheimer's disease, Parkinson's disease, Huntington's disease and multiple sclerosis, schizophrenia, psychosis or infantile hyperkinesia (ADHD, attention deficit / hyperactivity disorder) and other disorders mediated by the 5-HT₆ serotonin receptor in humans and/or in animals, preferably in mammals, more preferably in humans.

The preparation of the corresponding pharmaceutical compositions as well as the formulated medicaments can be carried out by means of conventional methods known in the prior art, for example, based on the indices of "Pharmaceutics: The Science of Dosage Forms", Second Edition, Aulton, M.E. (ED. Churchill Livingstone, Edinburgh (2002)); "Encyclopedia of Pharmaceutical Technology", Second Edition, Swarbrick, J. and Boylan, J.C. (Eds.), Marcel Dekker, Inc. New York (2002); "Modern Pharmaceutics", Fourth Edition, Banker G.S. and Rhodes C.T. (Eds.) Marcel Dekker, Inc. New York (2002), and "The Theory and Practice of Industrial Pharmacy", Lachman L., Lieberman H. and Kanig J. (Eds.), Lea & Febiger, Philadelphia (1986). The

respective literature descriptions are incorporated as a reference and are part of this disclosure.

The pharmaceutical compositions, as well as the formulated
5 medicaments prepared according to the present invention, can, in addition to at least one sulfonamide derivative of general formula (I), optionally in the form of one of its stereoisomers, preferably enantiomers or diastereomers, its racemate, or in form of a mixture of at least two of its stereoisomers, preferably enantiomers or diastereomers, in any mixing ratio, or a corresponding
10 physiologically acceptable salt thereof or a corresponding solvate thereof, comprise other conventional auxiliary substances known in the prior art, preferably excipients, fillers, solvents, diluents, dyes, coating agents, matrix forming agents and/or binders.

15 As the skilled persons in the art also knows, the choice of the auxiliary substances and the amounts thereof depend on the intended administration route, for example, rectal, intravenous, intraperitoneal, intramuscular, intranasal, oral, buccal or topical.

20 Medicaments suitable for oral administration are, for example, tablets, coated tablets, capsules or multiparticulates, preferably granules or pellets, optionally subjected to compression in tablets, filled in capsules or suspended in solutions, suspensions or suitable liquids.

25 Medicaments suitable for parenteral, topical or inhalatory administration may preferably be chosen from the group consisting of solutions, suspensions, quickly reconstitutable dry preparations and also sprays.

30 Medicaments suitable for oral or percutaneous use can release the sulfonamide compounds of general formula (I) in a sustained manner, the preparation of these sustained release medicaments generally being known in the prior art.

Suitable sustained release forms, as well as the materials and methods for the preparation thereof, are known in the prior art, for example from the indices of "Modified-Release Drug Delivery Technology", Rathbone, J.JI, Hadgraft, J. and Roberts, M.S. (Eds.), Marcel Dekker, Inc., New York (2002);
5 "Handbook of Pharmaceutical Controlled Release Technology", Wise, D.L. (Ed.), Marcel Dekker, Inc. New York (2000); "Controlled Drug Delivery", Vol. I, Basic Concepts, Bruck, S.D. (Ed.), CRD Press, Inc., Boca Raton (1983), and by Takada, K. and Yoshikawa, H., "Oral Drug Delivery", Encyclopedia of Controlled Drug Delivery, Mathiowitz, E. (Ed.), John Wiley & Sons, Inc., New York (1999),
10 Vol. 2, 728-742; Fix, J., "Oral drug delivery, small intestine and colon", Encyclopedia of Controlled Drug Delivery, Mathiowitz, E. (Ed.), John Wiley & Sons, Inc., New York (1999), Vol. 2, 698-728. The respective literature references are incorporated by reference and form part of the disclosure.

15 The medicament of the present invention may also have at least one enteric coating, which dissolves according to the pH. As a result of this coating, the medicament may pass through the stomach without dissolving, and the compounds of general formula I are only released in the intestinal tract. The enteric coating preferably dissolves at a pH of between 5 and 7.5. The materials
20 and methods suitable for preparing enteric coatings are also known in the prior art.

Typically, the pharmaceutical compositions and the medicaments comprise from 1 to 60% by weight of one or more sulfonamide derivatives of
25 general formula (I), and from 40 to 99% by weight of one or more excipients.

The medicament substance amount to be administered to the patient varies according to the patient's weight, the administration route, the indication and the severity of the disorder. Usually from 1 mg to 2 g of at least one
30 sulfonamide derivative of general formula (I) are administered per patient per day. The total daily dose can be administered to the patient in one or more doses.

Pharmaceutical Methods:

BINDING TO THE 5HT₆ SEROTONIN RECEPTOR

5

HEK-293 cell membranes expressing the recombinant human 5HT₆ receptor were supplied by Receptor Biology. The receptor concentration in said membranes is 2.18 pmol/mg of protein and the protein concentration is 9.17 mg/ml. The experimental protocol follows the method of B.L. Roth et al. [B.L. Roth, S.C. Craig, M.S. Choudhary, A. Uluer, F.J. Monsma, Y. Shen, H.Y. Meltzer, D.R. Sibley: Binding of Typical and Atypical Antipsychotic Agents to 5-Hydroxytryptamine-6 and Hydroxytryptamine-7 Receptors. *The Journal of Pharmacology and Experimental Therapeutics*, 1994, 268, 1403], with slight modifications. The commercial membrane is diluted (1:40 dilution) with the binding buffer: 50 mM Tris-HCl, 10 mM MgCl₂, 0.5 mM EDTA (pH 7.4). The radioligand used is [³H]-LSD at a concentration of 2.7 nM, the final volume being 200 µl. Incubation begins by adding 100 µl of the membrane suspension (≈ 22.9 µg of membrane protein), and is prolonged for 60 minutes at a temperature of 37°C. Incubation ends by quick filtration in a Harvester Brandel Cell through fiberglass filters of the Schleicher & Schuell GF 3362 trademark, pretreated with a 0.5% polyethyleneimine solution. The filters are washed three times with three milliliters of 50 mM Tris HCl buffer, pH 7.4. The filters are transferred to vials and 5 ml of Ecoscint H. liquid scintillation cocktail are added to each vial. The vials are left to equilibrate for several hours prior to their counting in a 1414 Wallac Winspectral scintillation counter. The non-specific binding is determined in the presence of 100 µM of serotonin. The assays are carried out in triplicate. The inhibition constants (K_i, nM) are calculated by non-linear regression analysis using the EBDA/LIGAND program [Munson and Rodbard, *Analytical Biochemistry*, 1980, 107, 220].

30

The respective literature descriptions are incorporated by reference and form part of the disclosure.

MEASUREMENTS OF FOOD INGESTION (BEHAVIOURAL MODEL)

Male W rats (200-270 g) from Harlan, S.A. are used. The animals are
5 acclimatized to the housings during at least 5 days prior to being subjected to
any treatment. During this period, the animals are housed (in groups of five) in
translucent cages and have free access to water and food. The animals are
housed in individual cages at least 24 hours prior to starting the treatment.

The acute effect of the sulfonamide derivatives of formula (I) used
10 inventively on food ingestion in rats in fasting conditions is then determined as
follows:

The rats are kept in fasting conditions for 23 hours in their individual
cages. After this period, the rats are orally or intraperitoneally treated with a
dose of a composition containing a sulfonamide derivative of general formula (I)
15 or a corresponding composition (vehicle) without said sulfonamide derivative.
Immediately after this, the rat is left with pre-weighed food and the accumulated
food intake is measured after 1, 2, 4 and 6 hours.

This food ingestion measuring method is also described in publications of
Kask et al., *European Journal of Pharmacology* 414 (2001), 215-224, and
20 Turnbull et al., *Diabetes*, Vol. 51, August, 2002. The respective bibliographic
descriptions are incorporated as a reference and they form part of the
disclosure.

The preparation of new compounds according to the invention is
indicated in the following examples. The affinity for the 5HT₆ serotonin receptor,
25 as well as the galenic formulas applicable to the compounds of the invention, is
also described. The examples indicated below, given as an illustrative example,
should in no way limit the scope of the invention.

Examples:

Example 1.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-5-chloro-3-methyl-benzo[b]thiophene-2-sulfonamide

5

185.6 mg (0.66 mMol) of 5-chloro-3-methyl-benzo[b] thiophene-2-sulfonyl chloride were added to a solution of 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole in 2 ml of dimethylformamide and 116 mg of N-ethyldiisopropylamine. The reaction mixture was stirred at the room temperature for 20 hours. Then it was evaporated to dryness, slightly alkalized with sodium bicarbonate solution and extracted with chloroform. The organic phase was repeatedly washed with water and saturated solution of sodium bicarbonate, it was separated and dried with anhydrous sodium sulfate. The organic solution was evaporated to dryness and the resulting solid was purified by chromatography, obtaining 180 mg (67%) of N-[1-(2-dimethylaminoethyl)-1H-indole-6-yl]-5-chloro-3-methyl-benzo[b]thiophene-2-sulfonamide as an amorphous solid.

10

15

Example 2.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-naphthalene-2-sulfonamide

20

187 mg (80%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 150 mg (0.66 mMol) of 2-naphthalenesulfonyl chloride, by means of the process described in the Example 1, as a solid.

25

Example 3.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-naphthalene-1-sulfonamide

157 mg (67%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 150 mg (0.66 mMol) of 1-naphthalenesulfonyl chloride, by means of the process described in the Example 1, as a solid.

Example 4.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-6-chloroimidazo[2,1-b]thiazole-5-sulfonamide

170 mg (67%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 170 mg (0.66 mMol) of 6-chloroimidazo[2,1-b]thiazole-5-sulfonyl chloride, by means of the process described in the Example 1, as a solid.

Example 5.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-4-phenylbenzenesulfonamide

184 mg (73%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 167 mg (0.66 mMol) of 4-phenylbenzenesulfonyl chloride, by means of the process described in the Example 1, as a solid.

Example 6.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-2-(naphthalene-1-yl)-ethanesulfonamide

100 mg (40%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 168 mg (0.66 mMol) of 2-(naphthalene-1-yl)-ethanesulfonyl chloride, by means of the process described in the Example 1, as a solid.

Example 7.- Preparation of N-[1-(2-Dimethylaminoethyl)-1H-indole-6-yl]-4-phenoxybenzenesulfonamide

190 mg (73%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 177 mg (0.66 mMol) of 4-phenoxybenzenesulfonyl chloride, by means of the process described in the Example 1, as a solid.

Example 8.- Preparation of N-[1-(2-dimethylaminoethyl)-1H-indole-6-yl]-3,5-dichlorobenzenesulfonamide

100 mg (41%) of the mentioned compound were obtained from 122 mg (0.6 mMol) of 6-amino-1-(2-dimethylaminoethyl)-1H-indole and 162 mg (0.66 mMol) of 3,5-dichlorobenzenesulfonyl chloride, by means of the process described in Example 1, as a solid.

Example 9.- Preparation of 5-Chloro-3-methyl-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-benzo[b]thiophene-2-sulfonamide

165 mg (58 %) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 186 mg (0.66 mMol) of 5-chloro-3-methyl-benzo[b]-thiophene-2-sulfonyl chloride by means of the process described in Example 1 as a solid.

Example 10.- Preparation of N-(1-[2-(Pyrrolidin-1-yl)ethyl]-1H-indol-6-yl)-naphthyl-2-sulfonamide

142 mg (57 %) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 150 mg (0.66 mMol) of naphthalenesulfonyl chloride by means of the process described in Example 1 as a solid.

Example 11.- Preparation of N-[1-[2-Pyrrolidin-1-yl]ethyl]-1H-indol-6-yl]-naphthalene-1-sulfonamide

166 (66 %) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 150 mg (0.66 mMol) naphthalenesulfonyl chloride by means of the process described in Example 1 as a solid.

Example 12.- Preparation of 6-Chloro-N-[1-[2-(pyrrolidin-1-yl)ethyl]-1H-indol-6-yl]-imidazo[2,1-b]thiazole-5-sulfonamide

170 mg (59%) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 170 mg 6-chloro-imidazo[2,1-b]thiazole-5-sulfonyl chloride by means of the process described in Example 1 as a solid.

Example 13.- Preparation of 4-Phenyl-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)benzenesulfonamide

205 mg (77%) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 169 mg (0.66 mmol) of 4-phenylbenzenesulfonyl chloride by means of the process described in Example 1 as an oil.

Example 14.- Preparation of 2-(Naphthyl-1-yl)-N-(1-(2-(pyrrolidin-1-yl)ethansulfonamid

182 mg (68%) of the mentioned compound were obtained from 137 mg (0.6 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 182 mg (0.66 mmol) of 2-naphthalene-ethansulfonyl chloride by means of the process described in Example 1 as a solid.

Example 15.- Preparation of 4-Phenoxy-N-(1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol-6-yl)-benzenesulfonamide


185 mg (67%) of the mentioned compound were obtained from 137 mg (0.6
5 mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 177 mg (0.66 mmol) of 4-phenoxybenzenesulfonyl chloride by means of the process described in Example 1 as a solid.

Example 16.-Preparation of 3,5-Dichloro-N-(1-(2-(Pyrrolidin-1-yl)-1H-indol-6-yl)-
10 benzenesulfonamide

122 mg (46%) of the mentioned compound were obtained from 137 mg (0.6
mMol) of 6-amino-1-(2-(pyrrolidin-1-yl)ethyl)-1H-indol and 162 mg (0.66 mmol) of 3,5-dichlorobenzenesulfonyl chloride by means of the process described in
15 Example 1 as a solid.




The yields are indicative and no added effort was made to improve them.




The melting point and spectroscopic data for identifying some of the
20 compounds of the present invention are indicated in the following table.



















Ex	R1	R2	R3	R4	R5	R6	R7	n	A	m.p. °C	IR cm ⁻¹	¹ H-NMR (300MHz), δ (solvent)
1	(CH ₃) ₂ N-	H	H	H	H	H	H	2		amorphous	3422, 3247, 2943, 1467, 1340, 1158, 1114, 1080, 862, 651,557.	2.19 (s, 9H); 2.55 (t, 2H, J=6.7 Hz); 4.13 (t, 2H, J=6.7 Hz); 6.42 (d, 1H, 3.1Hz); 6.69 (dd, 1H, J=8.3 Hz, J'=1.9 Hz); 7.13 (d, 1H, 3.1Hz); 7.23 (m, 1H); 7.45-7.37 (m, 2H); 7.63 (d, 1H, J=2.0 Hz); 7.69 (d, 1H, J=8.6 Hz). (DMSO-d6)
2	(CH ₃) ₂ N-	H	H	H	H	H	H	2		140-143	3422, 3246, 2935, 2760, 1468, 1336, 1159, 1132, 1074, 753, 711, 678, 553.	2.19 (s, 6H); 2.55 (t, 2H, J=7.0 Hz); 4.11 (t, 2H, J=7.0 Hz); 6.39 (d, 1H, J=3.1Hz); 6.67 (dd, 1H, J=8.3 Hz, J'=1.4 Hz); 7.10 (d, 1H, J=3.1Hz); 7.19 (s, 1H); 7.39 (d, 1H, J=8.4 Hz); 7.49-7.65 (m, 2H); 7.69 (dd, 1H, J=8.9 Hz, J'=1.6 Hz); 7.81-7.88 (m, 3H); 8.29 (s, 1H). (DMSO-d6)

PCT/EP 2004/008510

Ex	R1	R2	R3	R4	R5	R6	R7	n	A	m.p. °C	IR cm ⁻¹	¹ H-NMR (300MHz), δ (solvent)
3	(CH ₃) ₂ N-	H	H	H	H	H	H	2		139-142	3437, 2943, 1507, 1461, 1330, 1192, 1162, 1135, 961, 813, 763, 580, 472.	2.21 (s, 6H); 2.50 (t, 2H, J=7.0 Hz); 4.03 (t, 2H, J=7.0 Hz); 6.35 (d, 1H, J=3.1Hz); 6.48 (dd, 1H, J=8.4 Hz, J'=1.7 Hz); 7.00 (s, 1H); 7.05 (d, 1H, J=3.1Hz); 7.29 (m, 1H); 7.37 (t, 1H, J=7.8 Hz); 7.60 (m, 1H); 7.67 (m, 1H); 7.92 (d, 1H, J=8.1 Hz); 7.98 (d, 1H, J=8.1Hz); 8.10 (d, 1H, J=7.3 Hz); 8.73 (d, 1H, J=8.8 Hz). (DMSO-d6)
4	(CH ₃) ₂ N-	H	H	H	H	H	H	2		amorphous	3448, 3110, 2814, 1459, 1325, 1250, 1178, 1141, 621.	2.28 (s, 6H); 2.61 (t, 2H, J=7.0 Hz); 4.14 (t, 2H, J=7.0 Hz); 6.41 (d, 1H, J=3.1Hz); 6.81 (m, 2H); 7.12 (d, 1H, 3.1Hz); 7.19 (m, 1H); 7.42 (d, 1H, J=8.2 Hz); 7.56 (d, 1H, J=4.6 Hz) (DMSO-d6)
5	(CH ₃) ₂ N-	H	H	H	H	H	H	2		amorphous	3256, 2951, 2776, 1468, 1333, 1159, 1095, 763, 670, 591.	2.24 (s, 6H); 2.62 (t, 2H, J=7.0 Hz); 4.15 (t, 2H, J=7.0 Hz); 6.48 (d, 1H, J=3.1Hz); 6.70 (d, 1H, J=8.4 Hz); 7.12 (d, 1H, J=3.1Hz); 7.25 (d, 1H, J=3.3 Hz); 7.34-7.48 (m, 4H); 7.53 (m, 2H); 7.59 (AB sys, 2H, J=8.3 Hz); 7.78 (AB sys, 2H, J=8.3 Hz). (DMSO-d6)

Ex	R1	R2	R3	R4	R5	R6	R7	n	A	m.p. °C	IR cm ⁻¹	¹ H-NMR (300MHz), δ (solvent)
6	(CH ₃) ₂ N-	H	H	H	H	H	H	2		amorphous	3254, 2944, 1509, 1468, 1326, 1147, 970, 782, 716, 540.	2.15 (s, 6H); 2.62 (t, 2H, J=7.1 Hz); 3.38 (m, 2H); 3.49 (m, 2H); 4.22 (t, 2H, J=7.1 Hz); 6.47 (d, 1H, J=2.8Hz); 7.04 (m, 2H); 7.23 (d, 1H, J=3.1Hz); 7.26-7.45 (m, 5H); 7.56 (d, 1H, J=8.4 Hz); 7.68 (dd, 1H, J=7.5 Hz, J'=7.5 Hz); 7.77 (d, 1H, J=8.3Hz). (DMSO-d6)
7	(CH ₃) ₂ N-	H	H	H	H	H	H	2		amorphous	3255, 2935, 2768, 1583, 1488, 1334, 1245, 1154, 1093, 694, 570, 539.	2.28 (s, 6H); 2.65 (t, 2H, J=7.0 Hz); 4.16 (t, 2H, J=7.0 Hz); 6.42 (d, 1H, J=3.0Hz); 6.65 (dd, 1H, J=8.4 Hz, J'=1.7 Hz); 6.90 (AB sys, 2H, J=8.8 Hz); 7.00 (AB sys, 2H, J=7.9 Hz); 7.13 (d, 1H, J=3.1 Hz); 7.19 (m, 1H); 7.24 (m, 1H); 7.37 (m, 2H); 7.43 (d, 1H, J=8.3Hz); 7.65 (AB sys, 2H, J=8.9 Hz). (DMSO-d6)
8	(CH ₃) ₂ N-	H	H	H	H	H	H	2		150-159	3437, 3072, 2920, 1568, 1471, 1346, 1303, 1171, 1140, 799, 670, 598.	2.29 (s, 6H); 2.66 (t, 2H, J=6.8 Hz); 4.18 (t, 2H, J=6.8 Hz); 6.45 (d, 1H); 6.67 (d, 1H, J=8.4Hz); 7.15 (m, 1H); 7.19 (m, 1H); 7.46 (m, 2H); 7.59 (m, 2H). (DMSO-d6)

Ex	R1	R2	R3	R4	R5	R6	R7	n	A	p.f. °C	¹ H-RMN (300 MHz), δ (disolvante)
9		H	H	H	H	H	H	2		69-71	1,58(m, 4H); 2,31(m, 4H); 2,36(s, 3H); 2,59(m, 2H); 4,11(m, 2H); 6,31(s, 1H); 6,79(d, 1H, J=8,4 Hz); 7,09(s, 1H); 7,29(d, 1H, J=2,3 Hz); 7,38(d, 1H, J=8,5 Hz); 7,51(d, 1H, J=8,6 Hz); 7,94(d, 1H, J=1,0 Hz); 8,00(d, 1H, J=8,35 Hz); 10,39(b, 1H). (DMSO-d6)
10		H	H	H	H	H	H	2		54-60	1,54(m, 4H); 2,24(m, 4H); 2,50(m, 2H); 4,06(m, 2H); 6,25(s, 1H); 6,77(d, 1H, J=8,4 Hz); 7,07(s, 1H); 7,23(m, 1H); 7,32(d, 1H, J=8,1 Hz); 7,61(m, 2H); 7,75(d, 1H, J=8,8 Hz); 7,95(d, 1H, J=7,6 Hz); 8,03(m, 2H); 8,34(s, 1H); 10,11(b, 1H). (DMSO-d6)
11		H	H	H	H	H	H	2		160-165	1,74(m, 4H); 2,71(m, 4H); 2,94(m, 2H); 4,24(m, 2H); 6,27(d, 1H, J=2,8 Hz); 6,61(d, 1H, J=8,6 Hz); 7,09(s, 1H); 7,24(d, 1H, J=8,5 Hz); 7,28(d, 1H, J=2,8 Hz); 7,54(t, 1H, J=7,9 Hz); 7,63(m, 1H); 7,71(m, 1H); 8,03(d, 1H, J=7,6 Hz); 8,11-8,23(m, 2H); 8,77(d, 1H, J=8,2 Hz); 10,46(b, 1H). (DMSO-d6)
12		H	H	H	H	H	H	2		53-57	1,64(m, 4H); 2,50(m, 4H); 2,70(m, 2H); 4,14(m, 2H); 6,31(d, 1H, J=2,8 Hz); 6,71(d, 1H, J=8,8 Hz); 7,11(s, 1H); 7,31(d, 1H, J=2,9 Hz); 7,37(d, 1H, J=8,6 Hz); 7,56(d, 1H, J=4,4 Hz); 7,91(d, 1H, J=4,5 Hz); 10,63(b, 1H). (DMSO-d6)

13		H	H	H	H	H	H	H	2		178-181	1,55(m, 4H); 2,33(m, 4H); 2,61(m, 2H); 4,11(m, 2H); 6,30(d, 1H, J=2,8 Hz); 6,79(dd, 1H, J=8,2, J'=1,6 Hz); 7,09(s, 1H); 7,28(d, 1H, J=2,8 Hz); 7,34-7,49(m, 4H); 7,67(d, 2H, J=7,0 Hz); 7,76(AB sys, 2H, J=8,7 Hz); 7,80(AB sys, 2H, J=8,7 Hz); 10,05(bs, 1H). (DMSO-d6)
14		H	H	H	H	H	H	H	2		oil	1,49(m, 4H); 2,31(m, 4H); 2,66(t, 2H, J=6,5 Hz); 3,3 (m, 4H); 4,16(t, 2H, J=6,5 Hz); 6,40(dd, 1H, J=3,1 J'=0,7 Hz); 7,04(dd, 1H, J=8,4, J'=1,8 Hz); 7,13(m, 1H); 7,33-7,44(m, 5H); 7,48(d, 1H, J=8,6 Hz); 7,52(d, 1H, J=8,4 Hz); 7,75(t, 1H, J=4,8 Hz); 7,85(d, 1H, J=8,1 Hz); 9,84(s, 1H). (DMSO-d6)
15		H	H	H	H	H	H	H	2		59-62	1,61(m, 4H); 2,41(m, 4H); 2,66(t, 2H, J=6,5 Hz); 4,12(t, 2H, J=6,5 Hz); 6,30(d, 1H, J=2,8 Hz); 6,75(dd, 1H, J=8,4, J'=1,4 Hz); 6,99(d, 2H, J=8,8 Hz); 7,04(d, 2H, J=7,9 Hz); 7,10(s, 1H); 7,21(t, 1H, J=7,4 Hz); 7,29(d, 1H, J=3,1 Hz); 7,36(d, 1H, J=8,5 Hz); 7,41(t, 2H, J=7,9 Hz); 7,69(d, 2H, J=8,8 Hz); 9,98(bs, 1H). (DMSO-d6)
16		H	H	H	H	H	H	H	2		145-157	1,62(m, 4H); 2,39(m, 4H); 2,64(t, 2H, J=6,7 Hz); 4,15(t, 2H, J=6,7 Hz); 6,32(d, 1H, J=3,1 Hz); 6,73(dd, 1H, J=8,4, J'=1,8 Hz); 7,10(s, 1H); 7,33(d, 1H, J=3,2 Hz); 7,40(d, 1H, J=8,5 Hz); 7,63(d, 2H, J=1,9 Hz); 7,90(t, 1H, J=1,9 Hz); 10,20(bs, 1H). (DMSO-d6)

Pharmaceutical particulars:

Binding of the new compounds of general formula (I) to the 5-HT₆
 5 receptor was determined as previously described.

The binding results for some of the compounds of the present invention are indicated in the following table:

10	Table		
	Example	% Inhibition 10⁻⁶ M	K_i (nM)
	1	98.6	90.2
	2	97.7	41.2
15	3	95.3	19.8
	4	90.8	55.2
	5	93.4	129.4
	6	94.5	74.5
	7	95.1	118.6
20	8	86.9	159.1

The daily posology in human medicine is comprised between 1 milligram and 2 grams of medicinal product which can be administered in one or several doses. The compositions are prepared in forms that are compatible with the
 25 administration route used, preferably tablets, coated tablets, capsules, suppositories, solutions or suspensions. These compositions are prepared by means of known methods and comprise from 1 to 60% by weight of the medicament substance (compound of general formula I), and 40 to 99% by weight of the suitable pharmaceutical vehicle compatible with the medicament
 30 substance and the physical form of the composition used. The formula of a tablet containing a product of the invention is provided by way of example.

-66-

Example of formula per tablet:

Example 1	5 mg
Lactose	60 mg
Crystalline cellulose	25 mg
5 Povidone K 90	5 mg
Pregelatinized starch	3 mg
Colloidal silicon dioxide	1 mg
Magnesium stearate	1 mg
Total weight per tablet	100 mg